Oncology Clinical Pathways Kidney Cancer

October 2024 - V3.2024







Table of Contents

Presumptive Conditions.	3
Initial Workup.	4
Stage I	5
Stage II	6
Stage III.	7
Stage IV or Relapsed Clear Cell	8
<u>Surveillance</u>	9
Molecular Testing Table	10





<u>Kidney Cancer – Presumptive Conditions</u>

VA automatically presumes that certain disabilities were caused by military service. This is because of the unique circumstances of a specific Veteran's military service. If a presumed condition is diagnosed in a Veteran within a certain group, they can be awarded disability compensation.

Gulf War and Post 9/11 Veterans

If the patient served on or after Sept. 11, 2001, in Afghanistan, Djibouti, Egypt, Jordan, Lebanon, Syria, Uzbekistan, or Yemen or if the patient served in the *Southwest Asia theater of operations, or Somalia, on or after Aug. 2, 1990, specific conditions include:

Kidney cancer

*The Southwest Asia theater of operations refers to Iraq, Kuwait, Saudi Arabia, the neutral zone between Iraq and Saudi Arabia, Bahrain, Qatar, the United Arab Emirates, Oman, the Gulf of Aden, the Gulf of Oman, the Persian Gulf, the Arabian Sea, the Red Sea, and the airspace above these locations.


For more information, please visit <u>U.S. Department of Veterans Affairs - Presumptive Disability Benefits (va.gov)</u>







<u>Kidney Cancer – Initial Workup</u>



Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email CancerClinicalTrialsNavigation@va.gov.


- ^a Consider Genetic Counseling Evaluation for multiple or bilateral renal masses, family history, or young age
- ^b Consider Renal Biopsy if this would effect management
- ^c If Urothelial Carcinoma Suspected, consider urine cytology, ureteroscopy, or percutaneous biopsy
- d Treatment Decisions Dependent on patient comorbidities, surgical complexity, and shared decision making
- ^e If Clinically Indicated, consider bone scan, brain MRI, or chest CT







<u>Kidney Cancer – Stage I</u>



Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email CancerClinicalTrialsNavigation@va.gov.


- ^a Additional Options Include radical nephrectomy or ablative procedures including SBRT; renal biopsy; surveillance is preferred for small renal masses (≤3cm)
- ^b Active Surveillance appropriate in select patients (ex. not surgical candidates or medically frail)
- ^c Ablative Procedures including SBRT can be considered depending on case







Kidney Cancer - Stage II



Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email CancerClinicalTrialsNavigation@va.gov.

- ^a Nephrectomy Type dependent on location, comorbidities, and kidney function
- ^b Ablative Procedures including SBRT can be considered depending on case
- ^c **Pembrolizumab** consider if clear cell stage II with nuclear grade 4 or sarcomatoid differentiation, tumor stage III or higher, regional lymph-node metastasis, or stage M1 with NED, one year duration







<u>Kidney Cancer – Stage III</u>



Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email CancerClinicalTrialsNavigation@va.gov.


^a **Pembrolizumab** consider if clear cell stage II with nuclear grade 4 or sarcomatoid differentiation, tumor stage III or higher, regional lymph-node metastasis, or stage M1 with NED, one year duration







<u>Kidney Cancer – Stage IV or Relapsed Clear Cell</u>



Clinical trial(s) always considered on pathway. For assistance finding a clinical trial, email <u>CancerClinicalTrialsNavigation@va.gov</u>.

- ^a Intervention Includes either cytoreductive nephrectomy or metastasis directed therapy
- ^b **Prognostic Factors** include: less than one year from time of diagnosis to systemic therapy, performance status <80%, hemoglobin < lower limit of normal, calcium > upper limit of normal, neutrophil > upper limit of normal, and platelets > upper limit of normal
- ^c **Prognostic Risk Groups** defined as favorable (no prognostic factors), intermediate (1-2 prognostic factors), or poor (3-6 prognostic factors)
- d Pembrolizumab and Axitinib no ICI or TKI combination has shown overall survival in favorable risk; active surveillance may be used in certain situations

ICI Immune Checkpoint Inhibitor


TKI Tyrosine Kinase Inhibitor







<u>Kidney Cancer – Surveillance</u>









<u>Kidney Cancer – Molecular Testing Table</u>

Eligibility	Test Category	Test Type	Recommended Vendors	NPOP Coverage	Specimen Type
Kidney Cancer Stage 1-IV, Relapsed Clear Cell or Surveillance	No molecular testing is currently required for standard prognostication and therapy.				





Questions?

Contact VHAOncologyPathways@va.gov





